Volume 34/Number 12/December 1997

INDEX—*Volume 34*

Aguilar, A. Reports. Organochlorine pollutant levels in Mediterranean monk seals from the Western Mediterranean and the Sahara Coast 505
Aikman, P. Conference report 982
Albanis, T. A. Reports. Fungicide contamination of mediterranean estuarine waters: results from a MED Pol pilot survey 259
Albelmoschi, M. L. Reports. Correspondences between hydrology and particulate metal concentrations along mediterranean vertical profiles 479
Alden III, R. W. Reports. Optimizing temporal sampling strategies for benthic environmental monitoring programs 913
Al-Hadhrami, N. N. Reports. Studies on the biodegradation of three groups of pure n-alkanes in the presence of molasses and mineral fertiliser by pseudomonas aeruginosa 969
Allchin, C. R. Baseline. Metals and organochlorines in tissues of a Blainville's beaked whale (*mesoplodon densirostris*) and a killer whale (*orcinus orca*) stranded in the United Kingdom 208
Al-Lihaibi, S. S. Reports. Hydrocarbon distributions in sediemnts of the open area of the Arabian Gulf following the 1991 gulf war oil spill 941
Ambrose, P. Dead fish pollute bay 7
Ambrose, P. Latest El Nino warnings 855
Ambrose, P. News. 'Safer' sheep dips damage aquatic life 70
Ambrose, P. News. Break up of Russian tanker off Japan 146
Ambrose, P. News. Developments threaten unique marine habitats 854
Ambrose, P. News. Minamata Bay given the all clear 684
Ambrose, P. News. To dredge or not to dredge 223
Ambrose, P. News. Water hyacinth chokes Lake Victoria 364
Anderson, B. Viewpoint. General Guidelines for using the sediment quality triad 368
Andersson, R. Reports. Levels and patterns of polychlorinated dibenzo-*p*-dioxins, dibenzofurans and biphenyls in surface sediments from the western Baltic Sea (Arkona Basin) and the Oder River Estuarine System 1016
Anthony, R. G. Baseline. Organochlorines in sea otters and bald eagles from the Aleutian archipelago 486
Aral, O. Reports. Isomer-specific analysis of polychlorinated biphenyls in harbour porpose (phocoena phocoena) from the Black Sea 712
Aral, O. Reports. Persistent organochlorine residues in harbour porpoise (phocoena phocoena) from the Black Sea 338
Arnould, J. P. Y. Reports. Marine Debris Surveys at Bird Island, South Georgia 1990–1995 61
Asmund, G. Reports. Baseline levels and natural variability of element sin three seaweed species from West Greenland 171
Asmund, G. Reports. Uptake and release of lead and zinc by blue mussels. Experience from transplantion experiments in Greenland 805

Bacon, C. E. Baseline. Organochlorines in sea otters and bald eagles from the Aleutian archipelago 486
Baffi, F. Reports. Correspondences between hydrology and particulate metal concentrations along mediterranean vertical profiles 479
Baker, J. R. Baseline. Metals and organochlorines in tissues of a Blainville's beaked whale (*mesoplodon densirostris*) and a killer whale (*orcinus orca*) stranded in the United Kingdom 208
Balls, P. W. Reports. Physicochemical processes and their effects on the composition of suspended particulate material in estuaries: implications for monitoring and modelling 51
Balls, P. W. Reports. Trace metal in Scottish estuarine and coastal sediments 42
Ballschmiter, K. H. Viewpoint. Contamination of the deep-sea 288
Barcelo, D. Reports. Fungicide contamination of mediterranean estuarine waters: results from a MED Pol pilot survey 259
Bauer, B. Reports. Biological indicators used to map organotin contamination from a fishing port, Killybegs, Ireland 235
Bebianno, M. J. Baseline. Concentrations of metals and metallothioneins in *mytilus galloprovincialis* along the South Coast of Portugal 666
Bebianno, M. J. Reports. Sewage contamination following an accidental spillage in the Ria Formosa, Portugal 163
Beck, K. M. Baseline. Heavy metals in livers of bottlenose dolphins stranded along the south Carolina Coast 734

Beddig, S. Reports. A cadmium budget for the German Bight in the North Sea 375
Bellocq, J. Reports. Distribution and sources of polycyclic aromatic hydrocarbons in some Mediterranean coastal sediments 298
Bence, A. E. Baseline. An estimate of the annual input of natural petroleum hydrocarbons to seafloor sediments in Prince William Sound, Alaska 744
Benlahcen, K. T. Reports. Distribution and sources of polycyclic aromatic hydrocarbons in some Mediterranean coastal sediments 298
Bertrand, J. C. Reports. Hydrocarbon balance of a site which had been highly and chronically contaminated by petroleum wastes of a refinery (from 1956 to 1992) 456
Bester, K. Reports. Annual balances of hexachlorocyclohexanes, polychlorinated biphenyls and triazines in the German Bight 419
Blight, L. K. Reports. Occurrence of plastic particles in seabirds from the Eastern North Pacific 323
Boehm, P. D. Baseline. An estimate of the annual input of natural petroleum hydrocarbons to seafloor sediments in Prince William Sound, Alaska 744
Boireau, A. Reports. Polyaromatic hydrocarbon (PAH) distributions in the Seine river and its estuary 857
Bonfield, R. Reports. Distribution of technetium-99 in UK coastal waters 628
Borrell, A. Reports. Organochlorine pollutant levels in Mediterranean monk seals from the Western Mediterranean and the Sahara Coast 505
Boudou, A. Reports. Combined effects of water pH and salinity on the bioaccumulation of inorganic mercury and methylmercury in the shore crab carcinus maenas 880
Brockmann, U. Reports. A cadmium budget for the German Bight in the North Sea 375
Broman, D. Reports. Dioxin-like compounds in HPLC-fractionated extracts of marine samples from the east and west coast of Sweden: bioassay- and instrumentally-derived TCDD equivalents 1032
Broman, D. Reports. Impact of PAH outlets from an oil refinery on the receiving water area—sediment trap fluxes and multivariate statistical analysis 85
Brooks, R. Reports. Petroleum hydrocarbons in sediments of upper laguna madre 229
Brooks, T. Reports. Distribution of technetium-99 in UK coastal waters 628
Brown, J. Reports. Distribution of technetium-99 in UK coastal waters 628
Brugmann, L. Reports. Variability of dissolved and particulate trace metals in the Kiel and Mecklenburg Bights of the Baltic Sea 1990–1992 112
Bruni, V. Reports. Faecal pollution indicators in the Terra Nova Bay (Ross Sea, Antarctica) 908
Brunström, B. Reports. Dioxin-like compounds in HPLC-fractionated extracts of marine samples from the east and west coast of Sweden: bioassay- and instrumentally-derived TCDD equivalents 1032
Burger, A. E. Reports. Occurrence of plastic particles in seabirds from the Eastern North Pacific 323
Burns, W. A. Baseline. An estimate of the annual input of natural petroleum hydrocarbons to seafloor sediments in Prince William Sound, Alaska 744
Butler, E. C. V. Reports. Mercury in waters of the Derwent Estuary—Sample treatment and analysis 177
Butty, J. S. Baseline. Assessment of the octopus, octopus pallidus as a potential bioindicator species of xenobiotic challenge: baseline studies 564
Buxzinski, H. Reports. Distribution and sources of polycyclic aromatic hydrocarbons in some Mediterranean coastal sediments 298

Cabral, J. P. S. Reports. Biomonitoring with benthic macroalgae and direct assay of heavy metals in seawater of the Oporto coast (northwest Portugal) 1006
Caillet, G. Viewpoint. Contamination of the deep-sea 288
Cairns, Z. News. Ship specific tagging of oil contaminated discharges 70
Cameron, P. Baseline. Malformation rates in embryos of north sea fishes in 1991 and 1992 129
Cantillo, A. Y. Reports. Mollusc and sediment contaminant levels and trends in South Florida coastal waters 511
Carman, K. R. Reports. Effects of suspended diesel-contaminated sediment on feeding rate in the Darter Goby, gobionellus boleosoma (teleostei: Gobiidae) 269
Carr, S. Viewpoint. General Guidelines for using the sediment quality triad 368
Casini, S. Reports. Use of nondestructive biomarkers and residue analysis to access the health status of endangered species of pinnipeds in the South West Atlantic 157
Castelli, A. Analysis of macrozoobenthic community structure after severe dystrophic crises in a Mediterranean coastal lagoon 536

Castello, H. Reports. Use of nondestructive biomarkers and residue analysis to access the health status of endangered species of pinnipeds in the South West Altantic 157
Chan, C-C. Reports. Speciation study of chromium, copper and nickel in coastal estuarine sediments polluted by domestuic and industrial effluents 949
Chan, W-P. Reports. Speciation study of chromium, copper and nickel in coastal estuarine sediments polluted by domestuic and industrial effluents 949
Chaoui, A. Reports. Distribution and sources of polycyclic aromatic hydrocarbons in some Mediterranean coastal sediments 298
Chapman, P. Book review 981
Chapman, P. M. Editorial. Is bioaccumulation useful for predicting impacts 282
Chapman, P. News. OECD workshop recommends phasing out NOECs for aquatic toxicity test 145
Chapman, P. M. Correspondence. The Precautionary principle and ecological quality standards/objectives 227
Chapman, P. M. Editorial. Scientists need good manners to be scientists 852
Chapman, P. M. Viewpoint. General Guidelines for using the sediment quality triad 368
Chisholm, J. R. M. Reports. Wastewater discharge, seagrass decline and algal proliferation on the Cote d'Azur 78
Cini, R. Viewpoint. Adsorption and pollutant transport by marine aerosol 501
Clark, R. P. Reports. Organochlorines and other environmental contaminants in muscle tissues of sportfish collected from San Francisco Bay 1058
Cleveland, L. Reports. Toxicological and chemical screening of Antarctica sediments: use of whole sediment toxicity tests, microtox, mutatox and semipermeable membrane devices (SPMDs) 194
Cognetti, G. Conference Report. International marine congress in Palermo 675
Cognetti, G. Editorial. Rehabilitation and protection of the Black Sea: The challenge of a new transboundary environmental policy 752
Connor, M. S. Viewpoint. Tightening the relationship between monitoring and pollution abatement: the Massachusetts water resources authority contingency plan 9
Cook, J. M. Reports. The comparability of sample digestion techniques for the determination of metals in sediments 637
Correa, J. A. Reports. Epiphytic bacteria in copper-enriched environment in Northern Chile 816
Cote, W. A. Note. Potential effects of discarded triton paperboard six-pack carriers on fish and wildlife 135
Cotter, J. Reports. Distribution patterns of polychlorinated biphenyl congeners in water, sediment and biota from midway atoll (North Pacific Ocean) 548
Crockett, A. Reports. Toxicological and chemical screening of Antarctica sediments: use of whole sediment toxicity tests, microtox, mutatox and semipermeable membrane devices (SPMDs) 194
Croxall, J. P. Reports. Marine Debris Surveys at Bird Island, South Georgia 1990–1995 61
Cullen, M. Baseline. Marine debris on Northern New South Wales beaches (Australia): sources and the role of beach usage 348

dal Pont, G. Reports. Mercury in waters of the Derwent Estuary—Sample treatment and analysis 177
Dannecker, W. Reports. A cadmium budget for the German Bight in the North Sea 375
Dannenberger, D. Reports. Levels and patterns of polychlorinated dibenzo-*p*-dioxins, dibenzofurans and biphenyls in surface sediments from the western Baltic Sea (Arkona Basin) and the Oder River estuarine system 1016
Dassenakis, M. Reports. Trace metals transport and behaviour in the Mediterranean estuary of Acheloos River 103
Dauer, D. M. Reports. Optimizing temporal sampling strategies for benthic environmental monitoring programs 913
Davies, G. News. Offshore kazakhstan—ultimate challenge for the oil industry 145
Davies, M. Reports. Retroactive determination of industrial contaminations in tropical marine communities 975
Dawes, V. J. Reports. Polycyclic aromatic hydrocarbons (PAH) in seawater around England and Wales 306
Daza, J. L. Reports. Ecological analysis in a polluted area of Algeciras Bay (Southern Spain): External 'versus' internal outfalls and environmental implications 780
de Jonge, V. N. Reports. High remaining productivity in the Dutch Western Wadden Sea despite decreasing nutrient inputs from riverine sources 427
Dionne, J. Reports. Toxicological and chemical screening of Antarctica sediments: use of whole sediment toxicity tests, microtox, mutatox and semipermeable membrane devices (SPMDs) 194

Dixon, I. News. Toxic chemicals in great lakes to be eliminated 498
Donkin, M. E. Reports. Occurrence of marine antifouling agent irgarol 1051 within the Plymouth Sound locality: implications for the green macroalga enteromorpha intestinalis 645
Donkin, P. Reports. Occurrence of marine antifouling agent irgarol 1051 within the Plymouth Sound locality: implications for the green macroalga enteromorpha intestinalis 645
Douglas, G. S. Baseline. An estimate of the annual input of natural petroleum hydrocarbons to seafloor sediments in Prince William Sound, Alaska 744
Doumenq, P. Reports. Hydrocarbon balance of a site which had been highly and chronically contaminated by petroleum wastes of a refinery (from 1956 to 1992) 456
Doumenq, P. Reports. Long-term evolution of petroleum biomarkers in mangrove soil (Guadeloupe) 868
Downing, J. D. Reports. Organochlorines and other environmental contaminants in muscle tissues of sportfish collected from San Francisco Bay 1058
Ducrotoy, J-P. Viewpoint. Interrelations between science and policy-making: the North sea example 686
Duffield, D. A. Baseline. The determination by HPLC of PCB and p,p-DDE residues in marine mammals stranded on the Oregon coast, 1991–1995 844
Duggan, C. B. Reports. Biological indicators used to map organotin contamination from a fishing port, Killybegs, Ireland 235

Eertman, R. H. M. Reports. Toxicity of harbour canal sediments before dredging and after off-shore disposal 244
Elliott, A. J. Reports. The Influence of tidal currents and wind on mixing in the surface layers of the sea 15
Elliott, J. E. Reports. Heavy metal and metallothionein concentrations in seabirds from the Pacific coast of Canada 794
Elliott, M. Book review. Ecology of fjords and coastal waters 491
Elliott, M. Viewpoint. Interrelations between science and policy-making: the North sea example 686
Emmerson, R. H. C. Reports. A multivariate assessment of metal distribution in inter-tidal sediments of the Blackwater estuary, UK 960
Engle, V. Viewpoint. General Guidelines for using the sediment quality triad 368
Engwall, M. Reports. Dioxin-like compounds in HPLC-fractionated extracts of marine samples from the east and west coast of Sweden: bioassay- and instrumentally-derived TCDD equivalents 1032
Espinoza-Angulo, R. Reports. Fluxes and mass balances of nutrients in a semi-intensive shrimp farm in North-Western Mexico 290
Estacio, F. J. Reports. Ecological analysis in a polluted area of Algeciras Bay (Southern Spain): External 'versus' internal outfalls and environmental implications 780
Estes, J. A. Baseline. Organochlorines in sea otters and bald eagles from the Aleutian archipelago 486
Evans, S. M. Reports. Accumulation of beach litter on islands of the Pulau Seribu Archipelago, Indonesia 652

Fa, D. A. Reports. Ecological analysis in a polluted area of Algeciras Bay (Southern Spain): External 'versus' internal outfalls and environmental implications 780
Fair, P. Baseline. Heavy metals in livers of bottlenose dolphins stranded along the south Carolina Coast 734
Fairey, R. Reports. Organochlorines and other environmental contaminants in muscle tissues of sportfish collected from San Francisco Bay 1058
Falandysz, J. Baseline. Butyltin residues in sediment, fish, fish-eating birds, harbour porpoise and human tissues from the Polish Coast of the Baltic Sea 203
Featherstone, A. M. Reports. Removal of dissolved copper and iron at the freshwater-saltwater interface of an acid mine stream 332
Febvre, A. News. Oil spill research funded 682
Ferguson, A. Baseline. Metals, petroleum hydrocarbons and organochlorines in inshore sediments and waters of Mombassa, Kenya 570
Fernandez, M. B. Reports. Polyaromatic hydrocarbon (PAH) distributions in the Seine river and its estuary 857
Fernex, F. E. Reports. Wastewater discharge, seagrass decline and algal proliferation on the Cote d'Azur 78
Figueras, M. J. Viewpoint. Past, present and future perspectives on EU bathing water directive 148
Fileman, T. W. Reports. Occurrence of marine antifouling agent irgarol 1051 within the Plymouth Sound locality: implications for the green macroalga enteromorpha intestinalis 645

Fisher, P. J. Reports. Studies on the biodegradation of three groups of pure n-alkanes in the presence of molasses and mineral fertiliser by pseudomonas aeruginosa 969
Fleeger, J. W. Reports. Effects of suspended diesel-contaminated sediment on feeding rate in the Darter Goby, gobionellus boleosoma (teleostei: Gobiidae) 269
Flint, P. L. Reports. Persistence rates and detection probabilities of oiled king eider carcasses on St Paul Island, Alaska 522
Flores, V. Reports. Epiphytic bacteria in copper-enriched environment in Northern Chile 816
Fossi, M. C. Reports. Use of nondestructive biomarkers and residue analysis to access the health status of endangered species of pinnipeds in the South West Altantic 157
Fowler, A. C. Reports. Persistence rates and detection probabilities of oiled king eider carcasses on St Paul Island, Alaska 522
Frache, R. Reports. Correspondences between hydrology and particulate metal concentrations along mediterranean vertical profiles 479
Frances-Zubillaga, G. Reports. Enhanced growth in juvenile nereis diversicolor after its exposure to anaerobic polluted sediments 427
Frid, C. L. J. Reports. Abnormalities in the reproductive health of flounder platichthys flesus exposed to effluent from a sewage treatment works 34
Frid, C. L. J. Reports. The response of estuarine fish and benthos to an increasing discharge of sewage effluent 527
Froescheis, O. Viewpoint. Contamination of the deep-sea 288
Frost, A. Baseline. Marine debris on Northern New South Wales beaches (Australia): sources and the role of beach usage 348
Fryer, R. J. Reports. Designing monitoring programmes for detecting temporal trends in contaminants in fish and shellfish 821
Fukuda, M. Reports. Accumulation pattern of persistent organochlorine residues in common cormorants 186

Gabrielides, G. P. Reports. Fungicide contamination of mediterranean estuarine waters: results from a MED Pol pilot survey 259
Gaitis, A. Reports. Trace metals transport and behaviour in the Mediterranean estuary of Acheloos River 103
Galassi, S. Reports. Fungicide contamination of mediterranean estuarine waters: results from a MED Pol pilot survey 259
Galindo, R. J. G. Reports. Environmental and pollution condition of the Huizache-Caimanero Lagoon, in the north-west of Mexico 1072
Gangaiya, P. Baseline. Trace metal studies in the great Astrolabe Lagoon, Fiji, a pristine marine environment 353
Garcia-Adiego, E. M. Reports. Ecological analysis in a polluted area of Algeciras Bay (Southern Spain): External 'versus' internal outfalls and environmental implications 780
Garcia-gomez, J. C. Reports. Ecological analysis in a polluted area of Algeciras Bay (Southern Spain): External 'versus' internal outfalls and environmental implications 780
Gardner, M. J. Reports. The comparability of sample digestion techniques for the determination of metals in sediments 637
Garrigues, Ph. Reports. Distribution and sources of polycyclic aromatic hydrocarbons in some Mediterranean coastal sediments 298
Gerwinski, W. Reports. Lead budgets for the German Bight 410
Ghazi, S. J. Reports. Hydrocarbon distributions in sediemnts of the open area of the Arabian Gulf following the 1991 gulf war oil spill 941
Giddings, J. M. Editorial. Scientists need good manners to be scientists 852
Gill, M. E. Reports. Abnormalities in the reproductive health of flounder platichthys flesus exposed to effluent from a sewage treatment works 34
Gill, M. E. Reports. The response of estuarine fish and benthos to an increasing discharge of sewage effluent 527
Goldberg, E. D. Book Review. Tributylin 280
Gomez, E. Editorial. To save a reef 142
Gomez-Ariza, J. L. Reports. Ecological analysis in a polluted area of Algeciras Bay (Southern Spain): External 'versus' internal outfalls and environmental implications 780
Goodwin, M. News. Radioactive discharge controversy in France and Taiwan 754
Gracia, I. Reports. Speciation of heavy metals in sediments from salt marshes on the Southern Atlantic coast of Spain 123
Green, R. Viewpoint. General Guidelines for using the sediment quality triad 368
Gregg, J. C. Reports. Effects of suspended diesel-contaminated sediment on feeding rate in the Darter Goby, gobionellus boleosoma (teleostei: Gobiidae) 269
Griffiths, A. H. Reports. The comparability of sample digestion techniques for the determination of metals in sediments 637

Guard, M. Viewpoint. Dynamite fishing in Southern Tanzania, geographical variation, intensity of use and possible solutions 758
Guarro, J. Viewpoint. Past, present and future perspectives on EU bathing water directive 148
Guerrero-Galvan, S. R. Reports. Fluxes and mass balances of nutrients in a semi-intensive shrimp farm in North-Western Mexico 290
Guilano, M. Reports. Long-term evolution of petroleum biomarkers in mangrove soil (Guadeloupe) 868
Guiliano, M. Reports. Hydrocarbon balance of a site which had been highly and chronically contaminated by petroleum wastes of a refinery (from 1956 to 1992) 456
Guruge, K. S. Reports. Accumulation pattern of persistent organochlorine residues in common cormorants 186

Haarich, M. Reports. Lead budgets for the German Bight 410
Hall, J. A. Reports. The response of estuarine fish and benthos to an increasing discharge of sewage effluent 527
Hameedi, J. Viewpoint. General Guidelines for using the sediment quality triad 368
Hansen, H-P. Reports. Variability of dissolved and particulate trace metals in the Kiel and Mecklenburg Bights of the Baltic Sea 1990–1992 112
Harmon, M. Viewpoint. General Guidelines for using the sediment quality triad 368
Hart, M. A. Reports. Mercury accumulation in barred sand bass (paralabrax nebulifer) near a large wastewater outfall in the Southern Californian Bight 96
Haverland, P. Viewpoint. General Guidelines for using the sediment quality triad 368
Haynes, D. Baseline. Marine debris on continental islands and sand cays in the Far Northern Section of the Great Barrier Reef Marine Park, Australia 276
Haynes, D. Reports. A comparison of the bivalve species donax deltoides and mytilus edulis as monitors of metal exposure from effluent discharges along the ninety mile beach, Victoria, Australia 326
Hayteas, D. Baseline. The determination by HPLC of PCB and p,p-DDE residues in marine mammals stranded on the Oregon coast, 1991–1995 844
Heilprin, D. J. Reports. Mercury accumulation in barred sand bass (paralabrax nebulifer) near a large wastewater outfall in the Southern Californian Bight 96
Heinrich, H. Reports. Suspended particulate matter budget for the German Bight 398
Henderson, S. News. Bioremediation dechlorinator isolated 497
Henderson, S. News. Deliberate destruction of Californian Lake's ecosystem 365
Henderson, S. News. Drilling operations threaten Beaufort Sea whales 684
Henderson, S. News. Environmental agency action on oestrogenic discharges 284
Henderson, S. News. Future TBT phase-out likely 224
Henderson, S. News. How safe is dolphin-safe? 685
Henderson, S. News. Makah and Inuit to Resume Whaling 989
Henderson, S. News. Sea Turtles Under Threat 989
Hendrickx, M. E. Baseline. Distribution and abundance of meiofauna in a subtropical coastal lagoon on the south-eastern gulf of California, Mexico 582
Herd, R. A. Baseline. Metals, petroleum hydrocarbons and organochlorines in inshore sediments and waters of Mombassa, Kenya 570
Herut, B. Reports. Particulate metals contamination in the Kishon River Estuary, Israel 706
Hiscock, K. Viewpoint. Use available data 74
Holdway, D. A. Baseline. Assessment of the octopus, octopus pallidus as a potential bioindicator species of xenobiotic challenge: baseline studies 564
Hollyfield, S. Reports. Petroleum hydrocarbons in sediments of upper laguna madre 229
Holsbeek, L. Reports. Increase of organochlorines and mercury levels in common guillemots *Uria aalge* during winter in the southern North Sea 1049
Hope, B. Reports. Distribution patterns of polychlorinated biphenyl congeners in water, sediment and biota from midway atoll (North Pacific Ocean) 548
Hopkins, L. Oil and gas developments in the South Atlantic 286
Hortas, F. Reports. Ecological analysis in a polluted area of Algeciras Bay (Southern Spain): External 'versus' internal outfalls and environmental implications 780
Huggett, R. J. Reports. Assessment of Organotin in waters of selected gulf of Maine estuaries 802
Huhnerfuss, H. Reports. Annual balances of hexachlorocyclohexanes, polychlorinated biphenyls and triazines in the German Bight 419
Hull, S. Reports. Trace metal in Scottish estuarine and coastal sediments 42
Hyland, J. Viewpoint. General Guidelines for using the sediment quality triad 368

Ianni, C. Reports. Correspondences between hydrology and particulate metal concentrations along mediterranean vertical profiles 479
Ibarra, C. L. Reports. Environmental and pollution condition of the Huizache-Caimanero Lagoon, in the north-west of Mexico 1072
Ingersoll, C. Viewpoint. General Guidelines for using the sediment quality triad 368
Inza, I. Viewpoint. Past, present and future perspectives on EU bathing water directive 148
Izquierdo, C. Reports. Speciation of heavy metals in sediments from salt marshes on the Southern Atlantic coast of Spain 123

Jacquot, F. Reports. Hydrocarbon balance of a site which had been highly and chronically contaminated by petroleum wastes of a refinery (from 1956 to 1992) 456
Jacquot, F. Reports. Long-term evolution of petroleum biomarkers in mangrove soil (Guadeloupe) 868
Jarman, W. M. Baseline. Organochlorines in sea otters and bald eagles from the Aleutian archipelago 486
Jarman, W. M. Viewpoint. Contamination of the deep-sea 288
Jaubert, J. M. Reports. Wastewater discharge, seagrass decline and algal proliferation on the Cote d'Azur 78
Jepson, P. D. Baseline. Metals and organochlorines in tissues of a Blainville's beaked whale (mesoplodon densirostris) and a killer whale (orcinus orca) stranded in the United Kingdom 208
Jessep, M. A. Reports. The comparability of sample digestion techniques for the determination of metals in sediments 637
Johansen, P. Reports. Baseline levels and natural variability of element sin three seaweed species from West Greenland 171
Johansen, P. Reports. Uptake and release of lead and zinc by blue mussels. Experience from transplantion experiments in Greenland 805
Johnson, B. T. Reports. Toxicological and chemical screening of Antarctica sediments: use of whole sediment toxicity tests, microtox, mutatox and semipermeable membrane devices (SPMDs) 194
Johnson, E. Reports. Organochlorines and other environmental contaminants in muscle tissues of sportfish collected from San Francisco Bay 1058
Joiris, C. R. Reports. Increase of organochlorines and mercury levels in common guillemots *Uria aalge* during winter in the southern North Sea 1049
Jonas, P. J. C. Reports. Subduction and dispersion of a buoyant effluent plume in a stratified English bay 827
Jones, B. Reports. Distribution and speciation of heavy metals in surficial sediments from the Tees estuary, North-east England 768
Jones, B. R. Baseline. Metals and organochlorines in tissues of a Blainville's beaked whale (mesoplodon densirostris) and a killer whale (orcinus orca) stranded in the United Kingdom 208
Jones, P. News. Antarctic food chains disrupted? 597
Jones, P. News. Barriers taken off reef sanctuaries 71
Jones, P. News. India's coasts saved from shrimp peril 146
Jones, P. News. Poisoned fish walk on water 499
Jones, P. News. TBT implicated in mass dolphin deaths 146
Junin, M. Reports. Use of nondestructive biomarkers and residue analysis to access the health status of endangered species of pinnipeds in the South West Altantic 157

Kairu, K. K. Baseline. Metals, petroleum hydrocarbons and organochlorines in inshore sediments and waters of Mombassa, Kenya 570
Kan-Atireklap, S. Reports. Contamination by butyltin compounds in sediments from Thailand 894
Kannan, K. Baseline. Butyltin residues in sediment, fish, fish-eating birds, harbour porpoise and human tissues from the Polish Coast of the Baltic Sea 203
Kennedy, F. Reports. Effluent slicks in the Menai Strait: a comparison of ERS-1 Sar signatures and model predictions 264
Kim, E-Y. Baseline. Metal levels in some commercial fish species from Manila Bay, the Philippines 671
Korner, D. Reports. A cadmium budget for the German Bight in the North Sea 375
Kremling, K. Reports. Recent increase in the calcium concentrations in Baltic Sea Waters 763
Kremling, K. Reports. Variability of dissolved and particulate trace metals in the Kiel and Mecklenburg Bights of the Baltic Sea 1990–1992 112
Kress, N. Reports. Particulate metals contamination in the Kishon River Estuary, Israel 706

Kubilay, N. Baseline. The distribution of methultin species in different seas 739
Kubodera, T. Reports. Organochlorine residues in deep-sea organisms from Suruga Bay, Japan 250

Lam, M. H.-W. Reports. Speciation study of chromium, copper and nickel in coastal estuarine sediments polluted by domestuic and industrial effluents 949
Lamerdin, S. Reports. Organochlorines and other environmental contaminants in muscle tissues of sportfish collected from San Francisco Bay 1058
Landgraff, O. Reports. Annual balances of hexachlorocyclohexanes, polychlorinated biphenyls and triazines in the German Bight 419
Laporte, J. M. Reports. Combined effects of water pH and salinity on the bioaccumulation of inorganic mercury and methylmercury in the shore crab carcinus maenas 880
Lappin-Scott, H. M. Reports. Studies on the biodegradation of three groups of pure n-alkanes in the presence of molasses and mineral fertiliser by pseudomonas aeruginosa 969
Lardicci, C. Reports. Analysis of macrozoobenthic community structure after severe dystrophic crises in a Mediterranean coastal lagoon 536
Larsen, P. F. Reports. Assessment of Organotin in waters of selected gulf of Maine estuaries 802
Lasiak, T. Reports. Spatial and temporal variations in beach litter on the Transkei coast of South Africa 900
Laskaseru, B. O. Reports. Beach litter: an increasing and changing problem for Indonesia 469
Lauenstein, G. G. Reports. Mollusc and sediment contaminant levels and trends in South Florida coastal waters 511
Law, R. J. Baseline. Metals and organochlorines in tissues of a Blainville's beaked whale (*mesoplodon densirostris*) and a killer whale (*orcinus orca*) stranded in the United Kingdom 208
Law, R. J. Reports. Polycyclic aromatic hydrocarbons (PAH) in seawater around England and Wales 306
Leal, M. C. F. Reports. Biomonitoring with benthic macroalgae and direct assay of heavy metals in seawater of the Oporto coast (northwest Portugal) 1006
Le Dreau, Y. Reports. Hydrocarbon balance of a site which had been highly and chronically contaminated by petroleum wastes of a refinery (from 1956 to 1992) 456
Lebo, J. A. Reports. Toxicological and chemical screening of Antarctica sediments: use of whole sediment toxicity tests, microtox, mutatox and semipermeable membrane devices (SPMDs) 194
Lecke-Mitchell, K. M. Reports. Floating marine debris in the US Gulf of Mexico 702
Lee, J. S. Reports. Organochlorine residues in deep-sea organisms from Suruga Bay, Japan 250
Lee, R. F. Reports. Petroleum hydrocarbons and their effects in subtidal regions after major oil spills 928
Lee, W-S. Reports. Speciation study of chromium, copper and nickel in coastal estuarine sediments polluted by domestuic and industrial effluents 949
Leeder, J. Reports. A comparison of the bivalve species donax deltoides and mytilus edulis as monitors of metal exposure from effluent discharges along the ninety mile beach, Victoria, Australia 326
Leonard, K. S. Reports. Distribution of technetium-99 in UK coastal waters 628
Leonzio, C. Reports. Use of nondestructive biomarkers and residue analysis to access the health status of endangered species of pinnipeds in the South West Atlantic 157
Lester, J. N. Reports. A multivariate assessment of metal distribution in inter-tidal sediments of the Blackwater estuary, UK 960
Lewis, J. B. Reports. Abundance, distribution and partial mortality of the massive coral siderastrea siderea on degrading coral reefs at Barbados, West Indies 622
Little, E. E. Reports. Toxicological and chemical screening of Antarctica sediments: use of whole sediment toxicity tests, microtox, mutatox and semipermeable membrane devices (SPMDs) 194
Loglio, G. Viewpoint. Adsorption and pollutant transport by marine aerosol 501
Long, E. Viewpoint. General Guidelines for using the sediment quality triad 368
Long, E. R. Correspondence. On the identification of toxic hot spots using measures of the sediment quality triad 373
Lopes, C. F. Reports. Coastal monitoring program of Sao Sebastiao channel: assessing the effects of 'Tebar V' oil spill on rocky shore populations 923

Lorenzani, J. A. Reports. Use of nondestructive biomarkers and residue analysis to access the health status of endangered species of pinnipeds in the South West Atlantic 157
Lunel, T. Reports. The Influence of tidal currents and wind on mixing in the surface layers of the sea 15
Lye, C. M. Reports. Abnormalities in the reproductive health of flounder platichthys flesus exposed to effluent from a sewage treatment works 34

Machado, L. M. Baseline. Concentrations of metals and metallothioneins in mytilus galloprovincialis along the South Coast of Portugal 666
MacLeod, C. L. Reports. A multivariate assessment of metal distribution in inter-tidal sediments of the Blackwater estuary, UK 960
Madhusree, B. Reports. Isomer-specific analysis of polychlorinated biphenyls in harbour porpose (phocoena phocoena) from the Black Sea 712
Madhusree, B. Reports. Persistent organochlorine residues in harbour porpoise (phocoena phocoena) from the Black Sea 338
Madzena, A. Reports. Spatial and temporal variations in beach litter on the Transkei coast of South Africa 900
Mair, H. News. Fish kill reports along the eastern seaboard of America 756
Mair, H. News. Further leaks occur in sea empress saga 222
Mair, H. News. Intensive shrimp culture practices attacked by charity 6
Mair, H. News. North sea cod stocks close to collapse 144
Mair, H. News. Philippines tightens pollution controls 989
Mair, H. News. Sea Empress report published 682
Mair, H. News. Wildlife refuge experiment on midway atoll 366
Mankiewicz, P. J. Baseline. An estimate of the annual input of natural petroleum hydrocarbons to seafloor sediments in Prince William Sound, Alaska 744
Marsili, L. Reports. Use of nondestructive biomarkers and residue analysis to access the health status of endangered species of pinnipeds in the South West Altantic 157
Masaiganah, M. Viewpoint. Dynamite fishing in Southern Tanzania, geographical variation, intensity of use and possible solutions 758
Mathieu, D. Reports. Wastewater discharge, seagrass decline and algal proliferation on the Cote d'Azur 78
Matsuda, O. Reports. Comparative description of microbial community structure in surface sediments of eutrophic bays 26
Matthews, J. P. Reports. Effluent slicks in the Menai Strait: a comparison of ERS-1 Sar signatures and model predictions 264
Matthiessen, P. Reports. Polycyclic aromatic hydrocarbons (PAH) in seawater around England and Wales 306
Maugeri, T. L. Reports. Faecal pollution indicators in the Terra Nova Bay (Ross Sea, Antarctica) 908
Mayer, B. Reports. Suspended particulate matter budget for the German Bight 398
McCormick, D. Reports. Abnormalities in the reproductive health of flounder platichthys flesus exposed to effluent from a sewage treatment works 34
McCubbin, D. Reports. Distribution of technetium-99 in UK coastal waters 628
McFee, W. Baseline. Heavy metals in livers of bottlenose dolphins stranded along the south Carolina Coast 734
McIntyre, A. Book review 849
Medina, J. M. A. Reports. Environmental and pollution condition of the Huizache-Caimanero Lagoon, in the north-west of Mexico 1072
Mignotte, B. Reports. Comparative study of poliovirus recovery techniques from mussel tissues 875
Milanelli, J. C. C. Reports. Coastal monitoring program of Sao Sebastiao channel: assessing the effects of 'Tebar V' oil spill on rocky shore populations 923
Miles, A. K. Baseline. Organochlorines in sea otters and bald eagles from the Aleutian archipelago 486
Mille, G. Reports. Hydrocarbon balance of a site which had been highly and chronically contaminated by petroleum wastes of a refinery (from 1956 to 1992) 456
Mille, G. Reports. Long-term evolution of petroleum biomarkers in mangrove soil (Guadeloupe) 868
Miller, B. S. Reports. Trace metal in Scottish estuarine and coastal sediments 42
Minchin, D. Reports. Biological indicators used to map organotin contamination from a fishing port, Killybegs, Ireland 235
Miramand, P. Reports. Search for impact of a dredged material disposal site on growth and metal contamination of mytilus edulis (L.) in Charente-Maritime (France) 721

Miyazaki, N. Reports. Isomer-specific analysis of polychlorinated biphenyls in harbour porpose (phocoena phocoena) from the Black Sea 712
Miyazaki, N. Reports. Persistent organochlorine residues in harbour porpoise (phocoena phocoena) from the Black Sea 338
Monticelli, L. Reports. Faecal pollution indicators in the Terra Nova Bay (Ross Sea, Antarctica) 908
Morales, R. A. Reports. The Influence of tidal currents and wind on mixing in the surface layers of the sea 15
Moriki, A. Baseline. Dissolved trace elements in South Aegean seawater 840
Morrison, R. J. Baseline. Trace metal studies in the great Astrolabe Lagoon, Fiji, a pristine marine environment 353
Morton, B. Editorial. Asian Marine Research 494
Morton, B. Editorial. Hong Kong's First Marine Parks and Reserve 68
Mudge, S. Correspondence. Can vegetable oils outlast mineral oils in the marine environment? 213
Mudge, S. J. Reports. Sewage contamination following an accidental spillage in the Ria Formosa, Portugal 163
Mullin, K. Reports. Floating marine debris in the US Gulf of Mexico 702
Munoz, D. Reports. Long-term evolution of petroleum biomarkers in mangrove soil (Guadeloupe) 868

Näf, C. Reports. Dioxin-like compounds in HPLC-fractionated extracts of marine samples from the east and west coast of Sweden: bioassay- and instrumentally-derived TCDD equivalents 1032
Näf, C. Reports. Impact of PAH outlets from an oil refinery on the receiving water area—sediment trap fluxes and multivariate statistical analysis 85
Naidu, R. Baseline. Trace metal studies in the great Astrolabe Lagoon, Fiji, a pristine marine environment 353
Nakopoulou, C. Baseline. Dissolved trace elements in South Aegean seawater 840
Naqasima, M. R. Baseline. Trace metal studies in the great Astrolabe Lagoon, Fiji, a pristine marine environment 353
Newman, J. Reports. Organochlorines and other environmental contaminants in muscle tissues of sportfish collected from San Francisco Bay 1058
Nicholson, M. D. Reports. Designing monitoring programmes for detecting temporal trends in contaminants in fish and shellfish 821
Noguera, S. E. G. Baseline. Distribution and abundance of meiofauna in a subtropical coastal lagoon on the south-eastern gulf of California, Mexico 582
Norstrom, R. J. Baseline. Organochlorines in sea otters and bald eagles from the Aleutian archipelago 486
Nowell, D. News. Best practice in English coastal zone management 5

O'Connor, T. P. Reports. Mollusc and sediment contaminant levels and trends in South Florida coastal waters 511
O'Grady, B. V. Reports. Removal of dissolved copper and iron at the freshwater-saltwater interface of an acid mine stream 332
O'Reilly-Wiese, S. B. Reports. A multivariate assessment of metal distribution in inter-tidal sediments of the Blackwater estuary, UK 960
Ochoa-Acuña, H. Notes. Heavy metal concentrations in Juan Fernández fur seals (*arctocephalus philippii*) 663
Oehlmann, J. Reports. Biological indicators used to map organotin contamination from a fishing port, Killybegs, Ireland 235
Orazio, C. E. Reports. Toxicological and chemical screening of Antarctica sediments: use of whole sediment toxicity tests, microtox, mutatox and semipermeable membrane devices (SPMDs) 194
Owens, R. E. Reports. Physicochemical processes and their effects on the composition of suspended particulate material in estuaries: implications for monitoring and modelling 51
Ozdamar, E. Reports. Isomer-specific analysis of polychlorinated biphenyls in harbour porpose (phocoena phocoena) from the Black Sea 712
Ozdamar, E. Reports. Persistent organochlorine residues in harbour porpoise (phocoena phocoena) from the Black Sea 338
Ozturk, A. A. Reports. Persistent organochlorine residues in harbour porpoise (phocoena phocoena) from the Black Sea 338
Ozturk, A. A. Reports. Isomer-specific analysis of polychlorinated biphenyls in harbour porpose (phocoena phocoena) from the Black Sea 712
Ozturk, B. Reports. Isomer-specific analysis of polychlorinated biphenyls in harbour porpose (phocoena phocoena) from the Black Sea 712

Ozturk, B. Reports. Persistent organochlorine residues in harbour porpoise (phocoena phocoena) from the Black Sea 338

Paez-Osuna, F. Reports. Fluxes and mass balances of nutrients in a semi-intensive shrimp farm in North-Western Mexico 290
Page, D. S. Baseline. An estimate of the annual input of natural petroleum hydrocarbons to seafloor sediments in Prince William Sound, Alaska 744
Page, D. S. Reports. Petroleum hydrocarbons and their effects in subtidal regions after major oil spills 928
Pan, Y. Reports. Impacts of domestic sewage effluent on phytoplankton from Bedford Basin, eastern Canada 1001
Pastor, T. Reports. Organochlorine pollutant levels in Mediterranean monk seals from the Western Mediterranean and the Sahara Coast 505
Pearce, J. B. Book Review. Oil spills and other pollution 1078
Pearce, J. B. Book Review. Urbanization and marine ecosystems management 357
Pearce, J. Editorial. Marine pollution 592
Pearce, J. B. Book Review. Resolving scientific uncertainty 138
Pearce, N. J. G. Reports. Biomonitoring of pollution by *Cerastoderma edule* from the British Isles: a Laser Ablation ICP-MS study 1025
Petreas, M. Reports. Organochlorines and other environmental contaminants in muscle tissues of sportfish collected from San Francisco Bay 1058
Pettersen, H. Reports. Impact of PAH outlets from an oil refinery on the receiving water area—sediment trap fluxes and multivariate statistical analysis 85
Petty, J. D. Reports. Toxicological and chemical screening of Antarctica sediments: use of whole sediment toxicity tests, microtox, mutatox and semipermeable membrane devices (SPMDs) 194
Phillips, C. R. Reports. Mercury accumulation in barred sand bass (paralabrax nebulifer) near a large wastewater outfall in the Southern Californian Bight 96
Phillips, D. J. H. Editorial. Revisiting the southward tilt 362
Piatt, J. F. Reports. Mass-mortality of guillemots (uria aalge) in the Gulf of Alaska in 1993 656
Pirie, J. M. Reports. Trace metal in Scottish estuarine and coastal sediments 42
Plaschke, R. Reports. Mercury in waters of the Derwent Estuary—Sample treatment and analysis 177
Pohlmann, T. Reports. A cadmium budget for the German Bight in the North Sea 375
Pohlmann, T. Reports. Annual balances of hexachlorocyclohexanes, polychlorinated biphenyls and triazines in the German Bight 419
Polo, F. Viewpoint. Past, present and future perspectives on EU bathing water directive 148
Press, L. News. Conflicts of interest highlighted in Pacific Forum 754
Press, L. News. Lake Baikal seal deaths 496
Price, G. D. Reports. Biomonitoring of pollution by *Cerastoderma edule* from the British Isles: a Laser Ablation ICP-MS study 1025
Price, N. B. Reports. Physicochemical processes and their effects on the composition of suspended particulate material in estuaries: implications for monitoring and modelling 51
Proctor, W. Reports. Trace metal in Scottish estuarine and coastal sediments 42
Prosperi, V. A. Reports. Coastal monitoring program of Sao Sebastiao channel: assessing the effects of 'Tebar V' oil spill on rocky shore populations 923
Prudente, M. Baseline. Metal levels in some commercial fish species from Manila Bay, the Philippines 671
Puls, W. Reports. A cadmium budget for the German Bight in the North Sea 375
Puls, W. Reports. Lead budgets for the German Bight 410
Puls, W. Reports. Suspended particulate matter budget for the German Bight 398

Radach, G. Reports. A cadmium budget for the German Bight in the North Sea 375
Radach, G. Reports. Fluxes and budgets of contaminants in the German Bight 395
Radenac, G. Reports. Search for impact of a dredged material disposal site on growth and metal contamination of mytilus edulis (L.) in Charente-Maritime (France) 721
Rajemdran, R. Reports. Comparative description of microbial community structure in surface sediments of eutrophic bays 26
Rajendran, N. Reports. Comparative description of microbial community structure in surface sediments of eutrophic bays 26
Ranasinghe, J. A. Reports. Optimizing temporal sampling strategies for benthic environmental monitoring programs 913
Rao, D. V. S. Reports. Impacts of domestic sewage effluent on phytoplankton from Bedford Basin, eastern Canada 1001

Rappe, C. Reports. Levels and patterns of polychlorinated dibenzo-*p*-dioxins, dibenzofurans and biphenyls in surface sediments from the western Baltic Sea (Arkona Basin) and the Oder River estuarine system 1016
Ravenscroft, J. E. Reports. The comparability of sample digestion techniques for the determination of metals in sediments 637
Rayment, P. Reports. A comparison of the bivalve species donax deltoides and mytilus edulis as monitors of metal exposure from effluent discharges along the ninety mile beach, Victoria, Australia 326
Readman, J. W. Reports. Fungicide contamination of mediterranean estuarine waters: results from a MED Pol pilot survey 259
Rebers, A. Reports. A cadmium budget for the German Bight in the North Sea 375
Ree, T. News. First aid kit for oiled birds 365
Rees, I. News. SWISS- a new Irish Benthic survey 682
Rees, J. G. Baseline. Metals, petroleum hydrocarbons and organochlorines in inshore sediments and waters of Mombassa, Kenya 570
Reid, K. Reports. Marine Debris Surveys at Bird Island, South Georgia 1990–1995 61
Reid, P. C. Reports. Discharges from hydroelectric power schemes as a trigger for marine algal blooms 730
Review. Application of petroleum hydrocarbon chemical fingerprinting and allocation techniques after the Exxon Valdez Oil Spill 599
Rhudy, K. Reports. Petroleum hydrocarbons in sediments of upper laguna madre 229
Ribeyre, F. Reports. Combined effects of water pH and salinity on the bioaccumulation of inorganic mercury and methylmercury in the shore crab carcinus maenas 880
Rick, H-J. Reports. A cadmium budget for the German Bight in the North Sea 375
Riget, F. Reports. Baseline levels and natural variability of element sin three seaweed species from West Greenland 171
Riget, F. Reports. Uptake and release of lead and zinc by blue mussels. Experience from transplantion experiments in Greenland 805
Riquelme, C. Reports. Epiphytic bacteria in copper-enriched environment in Northern Chile 816
Ritterhoff, J. Reports. Influence of body length, life-history status and sex on trace metal concentrations in selected zooplankton collectives from the Greenland Sea 614
Rodgers, J. Jr: Viewpoint. General Guidelines for using the sediment quality triad 368
Rojas, A. Reports. Epiphytic bacteria in copper-enriched environment in Northern Chile 816
Roscigno, P. R. Reports. The effects of zinc contamination on the recruitment and early survival of benthic invertebrates in an estuary 443
Ross, C. A. Reports. Designing monitoring programmes for detecting temporal trends in contaminants in fish and shellfish 821
Rossi, F. Analysis of macrozoobenthic community structure after severe dystrophic crises in a Mediterranean coastal lagoon 536
Ruiz-Fernandez, A. C. Reports. Fluxes and mass balances of nutrients in a semi-intensive shrimp farm in North-Western Mexico 290

Saiz-Salinas, J. I. Reports. Enhanced growth in juvenile nereis diversicolor after its exposure to anaerobic polluted sediments 427
Salazar, M. Viewpoint. General Guidelines for using the sediment quality triad 368
Salihoglu, I. Baseline. The distribution of methultin species in different seas 739
Samsun, O. Reports. Isomer-specific analysis of polychlorinated biphenyls in harbour porpose (phocoena phocoena) from the Black Sea 712
Samsun, O. Reports. Persistent organochlorine residues in harbour porpoise (phocoena phocoena) from the Black Sea 338
Sangkoyo, H. Reports. Beach litter: an increasing and changing problem for Indonesia 469
Sanguansin, J. Reports. Contamination by butyltin compounds in sediments from Thailand 894
Savelli, C. Reports. Use of nondestructive biomarkers and residue analysis to access the health status of endangered species of pinnipeds in the South West Atlantic 157
Scarlett, A. Reports. Occurrence of marine antifouling agent irgarol 1051 within the Plymouth Sound locality: implications for the green macroalga enteromorpha intestinalis 645
Scatolini, S. Reports. Distribution patterns of polychlorinated biphenyl congeners in water, sediment and biota from midway atoll (North Pacific Ocean) 548
Schatzmann, M. Reports. A cadmium budget for the German Bight in the North Sea 375
Scherrer, P. Reports. Long-term evolution of petroleum biomarkers in mangrove soil (Guadeloupe) 868
Scheuhammer, A. M. Reports. Heavy metal and metallothionein concentrations in seabirds from the Pacific coast of Canada 794

Schirmacher, M. Reports. Lead budgets for the German Bight 410
Schlunzen, H. Reports. A cadmium budget for the German Bight in the North Sea 375
Schmidt, D. Reports. Lead budgets for the German Bight 410
Schulte-Oehlmann, U. Reports. Biological indicators used to map organotin contamination from a fishing port, Killybegs, Ireland 235
Schulz, M. Reports. A cadmium budget for the German Bight in the North Sea 375
Schwartzbrod, L. Reports. Comparative study of poliovirus recovery techniques from mussel tissues 875
Scott, P. J. B. Reports. Retroactive determination of industrial contaminations in tropical marine communities 975
Scoullos, M. Reports. Trace metals transport and behaviour in the Mediterranean estuary of Acheloos River 103
Selke, K. Reports. Annual balances of hexachlorocyclohexanes, polychlorinated biphenyls and triazines in the German Bight 419
Sepúlveda, M. S. Notes. Heavy metal concentrations in Juan Fernßndez fur seals (*arctocephalus philippii*) 663
Sharma, V. K. Reports. Petroleum hydrocarbons in sediments of upper laguna madre 229
Sharp, A. A. Reports. Sublethal effects of cupric ion activity on the grazing behaviour of three calanoid copepods 1041
Sharp, C. Reports. Subduction and dispersion of a buoyant effluent plume in a stratified English bay 827
Sheppard, S. Editorial. Biodiversity, productivity and system integrity 680
Sherwin, T. Reports. Subduction and dispersion of a buoyant effluent plume in a stratified English bay 827
Sherwin, T. J. Reports. Effluent slicks in the Menai Strait: a comparison of ERS-1 Sar signatures and model predictions 264
Sibley, P. K. Viewpoint. General Guidelines for using the sediment quality triad 368
Sicre, M-A. Reports. Polyaromatic hydrocarbon (PAH) distributions in the Seine river and its estuary 857
Sindermann, C. J. Editorial. The search for cause and effect relationships in marine pollution studies 218
Sindermann, C. Editorial. A marine pollution society 986
Smith, P. J. Viewpoint. General Guidelines for using the sediment quality triad 368
Sobral, P. Reports. Effects of copper exposure on the scope for growth of the clam *Ruditapes decussatus* from southern Portugal 992
Soggia, F. Reports. Correspondences between hydrology and particulate metal concentrations along mediterranean vertical profiles 479
Sommaripa, L. Viewpoint. Tightening the relationship between monitoring and pollution abatement: the Massachusetts water resources authority contingency plan 9
Sousa-Pinto, I. Reports. Biomonitoring with benthic macroalgae and direct assay of heavy metals in seawater of the Oporto coast (northwest Portugal) 1006
Spurrier, C. J. H. Baseline. Metals and organochlorines in tissues of a Blainville's beaked whale (*mesoplodon densirostris*) and a killer whale (*orcinus orca*) stranded in the United Kingdom 208
Stearns, D. E. Reports. Sublethal effects of cupric ion activity on the grazing behaviour of three calanoid copepods 1041
Stronkhorst, J. Reports. Toxicity of harbour canal sediments before dredging and after off-shore disposal 244
Sundermann, J. Reports. Fluxes and budgets of contaminants in the German Bight 395
Sundlof, S. F. Notes. Heavy metal concentrations in Juan Fernández fur seals (*arctocephalus philippii*) 663
Swartz, R. C. Viewpoint. General Guidelines for using the sediment quality triad 368

Taberski, K. Reports. Organochlorines and other environmental contaminants in muscle tissues of sportfish collected from San Francisco Bay 1058
Takemoto, N. Reports. Organochlorine residues in deep-sea organisms from Suruga Bay, Japan 250
Tan, K. S. Baseline. Imposex in three species of thais from Singapore, with additional observations on T. clavigera (kuster) from Japan 577
Tanabe, S. Baseline. Metal levels in some commercial fish species from Manila Bay, the Philippines 671
Tanabe, S. Reports. Accumulation pattern of persistent organochlorine residues in common cormorants 186
Tanabe, S. Reports. Contamination by butyltin compounds in sediments from Thailand 894

Tanabe, S. Reports. Isomer-specific analysis of polychlorinated biphenyls in harbour porpose (*phocoena phocoena*) from the Black Sea 712
Tanabe, S. Reports. Organochlorine residues in deep-sea organisms from Suruga Bay, Japan 250
Tanabe, S. Reports. Persistent organochlorine residues in harbour porpoise (*phocoena phocoena*) from the Black Sea 338
Tapia, G. Reports. Increase of organochlorines and mercury levels in common guillemots *Uria aalge* during winter in the southern North Sea 1049
Tardy, J. Reports. Search for impact of a dredged material disposal site on growth and metal contamination of mytilus edulis (L.) in Charente-Maritime (France) 721
Tatsukawa, R. Baseline. Metal levels in some commercial fish species from Manila Bay, the Philippines 671
Tatsukawa, R. Reports. Accumulation pattern of persistent organochlorine residues in common cormorants 186
Tatsukawa, R. Reports. Isomer-specific analysis of polychlorinated biphenyls in harbour porpose (*phocoena phocoena*) from the Black Sea 712
Tatsukawa, R. Reports. Persistent organochlorine residues in harbour porpoise (phocoena phocoena) from the Black Sea 338
Terver, D. Reports. Comparative study of poliovirus recovery techniques from mussel tissues 875
Thompson, B. Viewpoint. General Guidelines for using the sediment quality triad 368
Thompson, M. E. Note. Potential effects of discarded triton paperboard six-pack carriers on fish and wildlife 135
Tija, Y-W. Reports. Speciation study of chromium, copper and nickel in coastal estuarine sediments polluted by domestuic and industrial effluents 949
Titus, E. Reports. Distribution patterns of polychlorinated biphenyl congeners in water, sediment and biota from midway atoll (North Pacific Ocean) 548
Tokos, J. J. S. Reports. Variability of dissolved and particulate trace metals in the Kiel and Mecklenburg Bights of the Baltic Sea 1990–1992 112
Troncynski, J. Reports. Polyaromatic hydrocarbon (PAH) distributions in the Seine river and its estuary 857
Tronczynski, J. Reports. Fungicide contamination of mediterranean estuarine waters: results from a MED Pol pilot survey 259
Truchot, J. P. Reports. Combined effects of water pH and salinity on the bioaccumulation of inorganic mercury and methylmercury in the shore crab carcinus maenas 880
Truzzi, A. C. Reports. Coastal monitoring program of Sao Sebastiao channel: assessing the effects of 'Tebar V' oil spill on rocky shore populations 923
Tucci, S. Reports. Correspondences between hydrology and particulate metal concentrations along mediterranean vertical profiles 479
Tugrul, S. Baseline. The distribution of methultin species in different seas 739
Turki, A. Reports. Distribution and speciation of heavy metals in surficial sediments from the Tees estuary, North-east England 768

Uneputty, P. A. Reports. Accumulation of beach litter on islands of the Pulau Seribu Archipelago, Indonesia 652
Unger, M. A. Reports. Assessment of Organotin in waters of selected gulf of Maine estuaries 802
Urushigawa, Y. Reports. Comparative description of microbial community structure in surface sediments of eutrophic bays 26
Usero, J. Reports. Speciation of heavy metals in sediments from salt marshes on the Southern Atlantic coast of Spain 123

Van den Hurk, P. Reports. Toxicity of harbour canal sediments before dredging and after off-shore disposal 244
Van Pelt, T. I. Reports. Mass-mortality of guillemots (uria aalge) in the Gulf of Alaska in 1993 656
Varnavas, S. Baseline. Dissolved trace elements in South Aegean seawater 840
Vasconcelos, M. T. Reports. Biomonitoring with benthic macroalgae and direct assay of heavy metals in seawater of the Oporto coast (northwest Portugal) 1006
Vazquez, F. G. Reports. Petroleum hydrocarbons in sediments of upper laguna madre 229
Villagrana, L. C. Reports. Environmental and pollution condition of the Huizache-Caimanero Lagoon, in the north-west of Mexico 1072
Von Westernhagen, H. Baseline. Malformation rates in embryos of north sea fishes in 1991 and 1992 129
Voutsinou-Taliadouri, F. Baseline. Dissolved trace elements in South Aegean seawater 840

Walker, T. R. Reports. Marine Debris Surveys at Bird Island, South Georgia 1990–1995 61
Watzin, M. C. Reports. The effects of zinc contamination on the recruitment and early survival of benthic invertebrates in an estuary 443
Weisberg, S. B. Reports. Optimizing temporal sampling strategies for benthic environmental monitoring programs 913
Whitmarsh, D. Editorial. Artificial reefs: the law and the profits 2
Widdows, J. Reports. Effects of copper exposure on the scope for growth of the clam *Ruditapes decussatus* from southern Portugal 992
Wilhelm, G. Reports. Recent increase in the calcium concentrations in Baltic Sea Waters 763
Wilkins, T. News. IMO adopts draft guidelines on ballast water exchange 853
Wilkins, T. News. UK shifts its dumping stance 754
Williams, T. M. Baseline. Metals, petroleum hydrocarbons and organochlorines in inshore sediments and waters of Mombassa, Kenya 570
Willoughby, N. G. Reports. Beach litter: an increasing and changing problem for Indonesia 469
Wilson, C. J. Correspondence. On the identification of toxic hot spots using measures of the sediment quality triad 373
Windom, H. Viewpoint. General Guidelines for using the sediment quality triad 368
Wolf, D. Baseline. Heavy metals in livers of bottlenose dolphins stranded along the south Carolina Coast 734
Woodham, A. News. Snow goose population explosion may threaten Arctic ecosystems 990
Woodhead, R. J. Reports. Polycyclic aromatic hydrocarbons (PAH) in seawater around England and Wales 306
Yamagishi, S. Reports. Accumulation pattern of persistent organochlorine residues in common cormorants 186
Yates, R. Reports. The comparability of sample digestion techniques for the determination of metals in sediments 637
Yemenicioglu, S. Baseline. The distribution of methultin species in different seas 739
Yobe, A. C. Baseline. Metals, petroleum hydrocarbons and organochlorines in inshore sediments and waters of Mombassa, Kenya 570
Zanardi, E. Reports. Coastal monitoring program of Sao Sebastiao channel: assessing the effects of 'Tebar V' oil spill on rocky shore populations 923
Zauke, G-P. Reports. Influence of body length, life-history status and sex on trace metal concentrations in selected zooplankton collectives from the Greenland Sea 614
Zebühr, Y. Reports. Dioxin-like compounds in HPLC-fractionated extracts of marine samples from the east and west coast of Sweden: bioassay- and instrumentally-derived TCDD equivalents 1032